arXiv:solv-int/9901010v2 7 Feb 1999

February 9, 2008

Multi-soliton Solution of the Integrable Coupled Nonlinear Schrödinger Equation of Manakov Type

Freddy P. Zen[1)*] and Hendry I. Elim[1),2)†]
1) Theoretical High Energy Physics Group,
Theoretical Physics Lab., Department of Physics,
Institute of Technology Bandung, Bandung, Indonesia
2) Department of Physics, Pattimura University,
Ambon, Indonesia

Abstract

The general multi-soliton solution of the integrable coupled nonlinear Schrödinger equation (**NLS**) of Manakov type is investigated by using Zakharov-Shabat (**ZS**) scheme. We get the bright and dark multi-soliton solution using inverse scattering method of **ZS** scheme. Elastic and inelastic collision of N-solitons solution of the equation are also discussed.

PACS number(s): 42.65Sf

---

[*] email: fpzen@bdg.centrin.net.id
[†] email: hendry202@cyberlib.itb.ac.id

I. Introduction

The integrable coupled nonlinear Schrödinger equation of Manakov type is widely used in recent developments in the field of optical solitons in fibers. The use and applications of the equation is to explain how the solitons waves transmit in optical fiber, what happens when the interaction among optical solitons influences directly the capacity and quality of communication and so on.[1−4]

Some of exact solutions have been derived for the system related to the Manakov type equation with various methods.[5−10] Here, we will investigate the general solution of the following equation which is closely related to the Manakov type equation:

$$\begin{aligned} iq_{1_x} + \frac{(\alpha+\delta)}{\alpha(\alpha-\delta)} q_{1_{tt}} + 2\mu\left(|q_1|^2 + |q_2|^2\right) q_1 &= 0 \\ iq_{2_x} + \frac{(\alpha+\delta)}{\alpha(\alpha-\delta)} q_{2_{tt}} + 2\mu\left(|q_1|^2 + |q_2|^2\right) q_2 &= 0, \end{aligned} \tag{1.1}$$

where q_1 and q_2 are slowly varying envelopes of the two interacting optical modes, the variables x and t are the normalized distance and time, and α and δ are arbitrary real values which obey $\alpha \neq \delta$ and $\alpha \neq -\delta$. On the other hand, parameter μ is defined as follows

$$\mu = \pm\frac{\alpha^2 - \delta^2}{\alpha^2\delta}, \tag{1.2}$$

which is confirmed in section II.

Recently, in Ref.11, Radhakrishnan and Lakshmanan have derived the bright and dark multi-soliton solution of the related system in eq.(1.1) using Hirota method. However in this paper, we will investigate and provide another method to solve the related problem using Zakharov-Shabat inverse scattering method. We show that it is possible when the Zakharov-Shabat (**ZS**) scheme (in its final form) is expressed solely in terms of certain operators. Based on the choice of the operator, we find that there is a multi-soliton solution of eq.(1.1). We also find that there is an elastic and inelastic collision of the bright and dark multi-soliton.

This paper is organized as follows. In section II, we will perform the **ZS** scheme for eq.(1.1) and the arbitrary real parameter μ (eq.(1.2)). In section III, we will solve the bright and dark multi-soliton solution. We will compare the bright one and two soliton solution with the results in Ref.8 and Ref.10. Section IV is devoted for discussions and conclusions. In this section, we also state that there must be an integrable vector **NLS** system.

II. ZS Scheme for the Integrable Coupled NLS Equation of Manakov Type

We start by choosing the following two operators related to Zakharov-Shabat (**ZS**) scheme

$$\Delta_0^{(1)} = I\left(i\alpha\frac{\partial}{\partial x} - \frac{\partial^2}{\partial t^2}\right), \tag{2.1a}$$

and

$$\Delta_0^{(2)} = \left(\begin{array}{ccc} \alpha & 0 & 0 \\ 0 & \beta & 0 \\ 0 & 0 & \gamma \end{array}\right)\frac{\partial}{\partial t}, \tag{2.1b}$$

where α, β and γ are arbitrary real values, and I is the 3x3 unit matrix. We can then define the following operators by using this scheme related to inverse scattering techniques$^{12-14}$

$$\Delta^{(1)} = \Delta_0^{(1)} + U(t,x), \tag{2.2a}$$

and

$$\Delta^{(2)} = \Delta_0^{(2)} + V(t,x). \tag{2.2b}$$

Here operators $\Delta^{(i)}$,$(i=1,2)$ satisfy the following equation

$$\Delta^{(i)}(I + \Phi_+) = (I + \Phi_+)\Delta_0^{(i)}, \tag{2.3}$$

where the Volterra integral operators $\Phi_\pm(\psi)$ are defined according to equation

$$\Phi_\pm(\psi) = \int\limits_{-\infty}^{\infty} k_\pm(t,z)\,\psi(z)\,dz. \tag{2.4}$$

We now suppose that operators $\Phi_F(\psi)$ and $\Phi_\pm(\psi)$ are related to the following operator identity

$$(I + \Phi_+)(I + \Phi_F) = (I + \Phi_-), \tag{2.5}$$

where the integral operator $\Phi_F(\psi)$ is

$$\Phi_F(\psi) = \int_{-\infty}^{\infty} F(t,z)\,\psi(z)\,dz. \tag{2.6}$$

Both k_+ and F in eq.(2.4) and (2.6) are the 3x3 matrices chosen as follows

$$k_+ = \begin{pmatrix} a(t,z;x) & q_1(t,z;x) & q_2(t,z;x) \\ \pm q_1^*(t,z;x) & d(t,z;x) & e(t,z;x) \\ \pm q_2^*(t,z;x) & f(t,z;x) & g(t,z;x) \end{pmatrix}, \tag{2.7a}$$

and

$$F = \begin{pmatrix} 0 & A_n(t,z;x) & B_n(t,z;x) \\ A_n^*(t,z;x) & 0 & 0 \\ B_n^*(t,z;x) & 0 & 0 \end{pmatrix}. \tag{2.7b}$$

Here a, q_1, q_2, $\pm q_1^*$, $\pm q_2^*$, d, e, f, g, A_n, A_n^*, B_n and B_n^* are parameters which will be calculated in section III.

In eq.(2.5), we have assumed that $(I + \Phi_+)$ is invertible, then

$$(I + \Phi_F) = (I + \Phi_+)^{-1}(I + \Phi_-), \tag{2.8}$$

so that operator $(I + \Phi_F)$ is factorisable. From eq.(2.5), we derive Marchenko matrix equations,

$$k_+(t,z;x) + F(t,z;x) + \int_t^{\infty} k_+(t,t';x)\,F(t',z;x)\,dt' = 0, \quad \text{for } z > t\,, \tag{2.9a}$$

and

$$k_-(t,z;x) = F(t,z;x) + \int_t^{\infty} k_+(t,t';x)\,F(t',z;x)\,dt'\,, \quad \text{for } z < t. \tag{2.9b}$$

In eq.$(2.9b)$, k_- is obviously defined in terms of k_+ and F. Both eq.$(2.9a)$ and $(2.9b)$ require F, and F is supplied by the solution of equations :

$$\left[\Delta_0^{(1)}, \Phi_F\right] = 0, \tag{2.10a}$$

and

$$\left[\Delta_0^{(2)}, \Phi_F\right] = 0. \tag{2.10b}$$

After a little algebraic manipulation, we get

$$i\alpha F_x + F_{zz} - F_{tt} = 0, \tag{2.11a}$$

and

$$\begin{pmatrix} \alpha & 0 & 0 \\ 0 & \beta & 0 \\ 0 & 0 & \gamma \end{pmatrix} F_t + F_z \begin{pmatrix} \alpha & 0 & 0 \\ 0 & \beta & 0 \\ 0 & 0 & \gamma \end{pmatrix} = 0, \tag{2.11b}$$

where $F_t \equiv \frac{\partial F}{\partial t}$, $F_z \equiv \frac{\partial F}{\partial z}$, etc.

$U\left(t,x\right)$ and $V\left(t,x\right)$ can be found by solving eq.(2.3) in which we have substituted eq.(2.7a) and (2.4) (for k_+) to that equation:

$$V\left(t,x\right) = \begin{pmatrix} 0 & \left(\alpha-\beta\right)q_1 & \left(\alpha-\gamma\right)q_2 \\ \pm\left(\beta-\alpha\right)q_1^* & 0 & \left(\beta-\gamma\right)e \\ \pm\left(\gamma-\alpha\right)q_2^* & \left(\gamma-\beta\right)f & 0 \end{pmatrix}, \tag{2.12a}$$

and

$$U\left(t,x\right) = -2k_{+_t} = -2\begin{pmatrix} a_t & q_{1_t} & q_{2_t} \\ \pm q_{1_t}^* & d_t & e_t \\ \pm q_{2_t}^* & f_t & g_t \end{pmatrix}. \tag{2.12b}$$

Based on the solution of equation $\Delta^{(2)}\left(I + \Phi_+\right) = \left(I + \Phi_+\right)\Delta_0^{(2)}$, we get that $k_+\left(t,z;x\right)$ must obey the following equation:

$$\begin{pmatrix} \alpha & 0 & 0 \\ 0 & \beta & 0 \\ 0 & 0 & \gamma \end{pmatrix} k_{+t} + k_{+z}\begin{pmatrix} \alpha & 0 & 0 \\ 0 & \beta & 0 \\ 0 & 0 & \gamma \end{pmatrix}$$

$$+\begin{pmatrix} 0 & \left(\alpha-\beta\right)q_1 & \left(\alpha-\gamma\right)q_2 \\ \pm\left(\beta-\alpha\right)q_1^* & 0 & \left(\beta-\gamma\right)e \\ \pm\left(\gamma-\alpha\right)q_2^* & \left(\gamma-\beta\right)f & 0 \end{pmatrix} k_+ = 0 \tag{2.13}$$

and if we evaluate this eq.(2.13) on $z = t$, we find

$$a_t = \mp\frac{1}{\alpha}\left[\left(\alpha-\beta\right) \mid q_1 \mid^2 + \left(\alpha-\gamma\right)\left|q_2\right|^2\right], \tag{2.14a}$$

$$d_t = -\frac{1}{\beta}\left[\left(\beta-\gamma\right)ef \pm \left(\beta-\alpha\right)\left|q_1\right|^2\right], \tag{2.14b}$$

$$e_t = -\frac{1}{\beta}\left[(\beta-\gamma)\,eg \pm (\beta-\alpha)\,q_1^* q_2\right], \tag{2.14c}$$

$$f_t = -\frac{1}{\gamma}\left[(\gamma-\beta)\,fd \pm (\gamma-\alpha)\,q_2^* q_1\right], \tag{2.14d}$$

$$g_t = -\frac{1}{\gamma}\left[(\gamma-\beta)\,ef \pm (\gamma-\alpha)\,|q_2|^2\right]. \tag{2.14e}$$

After substituting the above equations into eq.$(2.12b)$ and eq.$(2.2a)$, we find the following equation

$$\Delta^{(1)} = I\left(i\alpha\frac{\partial}{\partial x} - \frac{\partial^2}{\partial t^2}\right) - 2\begin{pmatrix} a_t & q_{1_t} & q_{2_t} \\ \pm q_{1_t}^* & d_t & e_t \\ \pm q_{2_t}^* & f_t & g_t \end{pmatrix}. \tag{2.15}$$

On the other hand, by substituting eq.$(2.12a)$ into eq. $(2.2b)$, we get

$$\Delta^{(2)} = \begin{pmatrix} \alpha & 0 & 0 \\ 0 & \beta & 0 \\ 0 & 0 & \gamma \end{pmatrix}\frac{\partial}{\partial t} + \begin{pmatrix} 0 & (\alpha-\beta)\,q_1 & (\alpha-\gamma)\,q_2 \\ \pm(\beta-\alpha)\,q_1^* & 0 & (\beta-\gamma)\,e \\ \pm(\gamma-\alpha)\,q_2^* & (\gamma-\beta)\,f & 0 \end{pmatrix}. \tag{2.16}$$

Since $\Delta^{(1)}$ commutes with $\Delta^{(2)}$,

$$\begin{aligned} iq_{1_x} + \frac{(\alpha+\beta)}{\alpha(\alpha-\beta)}q_{1_{tt}} \pm 2\left(\frac{1}{\alpha(\alpha-\beta)}\right)\left(\left[\theta_1 \mid q_1 \mid^2 + \theta_2 |q_2|^2\right] q_1 + Q_1\right) &= 0 \\ iq_{2_x} + \frac{(\alpha+\beta)}{\alpha(\alpha-\beta)}q_{2_{tt}} \pm 2\left(\frac{1}{\alpha(\alpha-\beta)}\right)\left(\left[\theta_3 \mid q_1 \mid^2 + \theta_4 |q_2|^2\right] q_2 + Q_2\right) &= 0, \end{aligned} \tag{2.17a}$$

and its complex conjugate

$$\begin{aligned} -iq_{1_x}^* + \frac{(\alpha+\beta)}{\alpha(\alpha-\beta)}q_{1_{tt}}^* \pm \frac{2}{\alpha(\alpha-\beta)}\left(\left[\theta_5 \mid q_1 \mid^2 + \theta_6 |q_2|^2\right] q_1^* + Q_1^*\right) &= 0 \\ -iq_{2_x}^* + \frac{(\alpha+\beta)}{\alpha(\alpha-\beta)}q_{2_{tt}}^* \pm \frac{2}{\alpha(\alpha-\beta)}\left(\left[\theta_7 \mid q_1 \mid^2 + \theta_8 |q_2|^2\right] q_2^* + Q_2^*\right) &= 0, \end{aligned} \tag{2.17b}$$

where

$$\theta_1 = \theta_5 = \frac{1}{\alpha\beta}\left[(\alpha+\beta)(\alpha-\beta)^2\right], \tag{2.18a}$$

$$\theta_2 = \frac{1}{\alpha\gamma}\left[\gamma(\alpha-\beta)(\alpha-\gamma)+\alpha(\alpha-\gamma)^2\right], \tag{2.18b}$$

$$\theta_3 = \frac{1}{\alpha\beta}\left[\beta(\alpha-\beta)(\alpha-\gamma)+\alpha(\alpha-\beta)^2\right], \tag{2.18c}$$

$$\theta_4 = \theta_8 = \frac{1}{\alpha\gamma}\left[(\alpha+\gamma)(\alpha-\gamma)^2\right], \tag{2.18d}$$

$$\theta_6 = \frac{1}{\alpha\beta}\left[(\alpha+\beta)(\alpha-\beta)(\alpha-\gamma)\right], \tag{2.18e}$$

$$\theta_7 = \frac{1}{\alpha\gamma}\left[(\alpha+\gamma)(\alpha-\beta)(\alpha-\gamma)\right], \tag{2.18f}$$

$$Q_1 = \left[\frac{(\alpha-\beta)(\beta-\gamma)}{\beta}efq_1+(\gamma-\beta)fq_{2_t}-\frac{(\alpha-\gamma)(\gamma-\beta)}{\gamma}fdq_2\right], \tag{2.18g}$$

$$Q_2 = \left[\frac{(\alpha-\gamma)(\gamma-\beta)}{\gamma}efq_2+(\beta-\gamma)eq_{1_t}-\frac{(\alpha-\beta)(\beta-\gamma)}{\beta}egq_1\right], \tag{2.18h}$$

$$e = f^* \text{ and } d = \frac{\gamma}{\beta}g^*. \tag{2.18i}$$

If we put parameters $\beta=\gamma=\delta$ (where δ is an arbitrary real parameter) into the above equations and then substitute the results into eq.$(2.17a)$ and $(2.17b)$, we find

$$\begin{aligned} iq_{1_x} + \frac{(\alpha+\delta)}{\alpha(\alpha-\delta)}q_{1_{tt}} + 2\mu\left[|q_1|^2+|q_2|^2\right]q_1 &= 0 \\ i\,q_{2_x} + \frac{(\alpha+\delta)}{\alpha(\alpha-\delta)}q_{2_{tt}} + 2\mu\left[|q_1|^2+|q_2|^2\right]q_2 &= 0, \end{aligned} \tag{2.19a}$$

and its complex conjugate

$$\begin{aligned} -iq^*_{1_x} + \frac{(\alpha+\delta)}{\alpha(\alpha-\delta)}q^*_{1_{tt}} + 2\mu\left[|q_1|^2+|q_2|^2\right]q^*_1 &= 0 \\ -iq^*_{2_x} + \frac{(\alpha+\delta)}{\alpha(\alpha-\delta)}q^*_{2_{tt}} + 2\mu\left[|q_1|^2+|q_2|^2\right]q^*_2 &= 0. \end{aligned} \tag{2.19b}$$

Here parameter μ is an arbitrary real value which has been defined in eq.(1.2). It is obvious that eq.(2.19a) and (2.19b) are the general form of the integrable coupled nonlinear Schrödinger equation of Manakov type (eq.(1.1)) and its complex conjugate. We note that the functions Q_1 and Q_2 in eq.(2.17a) and its complex conjugates in eq.(2.17b) are the perturbative terms (inhomogeneous terms) for arbitrary α, β and γ.

III. The Bright and Dark Multi-Soliton Solution

We consider a general matrix function F in eq.(2.7b) and substitute it into eq.(2.11a), we find four differential equations

$$\begin{aligned} i\alpha A_{n_x} + A_{n_{zz}} - A_{n_{tt}} &= 0, && (3.1a)\\ i\alpha B_{n_x} + B_{n_{zz}} - B_{n_{tt}} &= 0, && (3.1b)\\ i\alpha A^*_{n_x} + A^*_{n_{zz}} - A^*_{n_{tt}} &= 0, && (3.1c)\\ i\alpha B^*_{n_x} + B^*_{n_{zz}} - B^*_{n_{tt}} &= 0\,. && (3.1d) \end{aligned}$$

The solution of the above equations can be derived by using separable variable method. We then find

$$\begin{aligned} A_n\left(t,z;x\right) &= \sum_{n=1}^{N} A_{n_0} e^{-\alpha\rho_n z}\left[e^{\rho_n\left(\delta t + i\rho_n\left(\alpha^2-\delta^2\right)x\right)}\right], && (3.2a)\\ B_n\left(t,z;x\right) &= \sum_{n=1}^{N} B_{n_0} e^{-\alpha\rho_n z}\left[e^{\rho_n\left(\delta t + i\rho_n\left(\alpha^2-\delta^2\right)x\right)}\right], && (3.2b)\\ A^*_n\left(t,z;x\right) &= \sum_{n=1}^{N} A^*_{n_0} e^{-\delta\sigma_n z}\left[e^{\sigma_n\left(\alpha t + i\sigma_n\left(\delta^2-\alpha^2\right)x\right)}\right], && (3.2c)\\ B^*_n\left(t,z;x\right) &= \sum_{n=1}^{N} B^*_{n_0} e^{-\delta\sigma_n z}\left[e^{\sigma_n\left(\alpha t + i\sigma_n\left(\delta^2-\alpha^2\right)x\right)}\right], && (3.2d) \end{aligned}$$

where A_{n_0} , $A^*_{n_0}$, B_{n_0} and $B^*_{n_0}$ are arbitrary complex parameters.

To get the final solution of the integrable coupled **NLS** equation of Manakov type, we have to substitute the equations (2.7b), (3.2a), (3.2b), (3.2c) and (3.2d) into Marchenko matrix equation (eq.(2.9a)). We get (for a, q_1, and q_2)

$$
\begin{aligned}
a \;=\; & -\sum_{n=1}^{N} \int_{t}^{\infty} q_1\, A_{n_0}^{*} e^{i\frac{\sigma_n^2}{\alpha}\left(\delta^2-\alpha^2\right)x} e^{-\delta\sigma_n z} e^{\alpha\sigma_n t'} dt' \\
& \qquad -\sum_{n=1}^{N} \int_{t}^{\infty} q_2 B_{n_0}^{*} e^{i\frac{\sigma_n^2}{\alpha}\left(\delta^2-\alpha^2\right)x} e^{-\delta\sigma_n z} e^{\alpha\sigma_n t'} dt', \qquad (3.3\text{a})
\end{aligned}
$$

$$
\begin{aligned}
q_1 \;=\; & -\sum_{n=1}^{N} \left(e^{-\alpha\rho_n z} A_{n_0}\right) e^{\delta\rho_n t} e^{i\frac{\rho_n^2}{\alpha}\left(\alpha^2-\delta^2\right)x} \\
& -\sum_{n=1}^{N} \left(A_{n_0}\right) \int_{t}^{\infty} a\left(t,z;x\right)\, e^{i\frac{\rho_n^2}{\alpha}\left(\alpha^2-\delta^2\right)x} e^{-\alpha\rho_n z} e^{\delta\rho_n t'} dt'\,, \qquad (3.3\text{b})
\end{aligned}
$$

and

$$
\begin{aligned}
q_2 \;=\; & -\sum_{n=1}^{N} \left(e^{-\alpha\rho_n z} B_{n_0}\right) e^{\delta\rho_n t} e^{i\frac{\rho_n^2}{\alpha}\left(\alpha^2-\delta^2\right)x} \\
& -\sum_{n=1}^{N} \left(B_{n_0}\right) \int_{t}^{\infty} a\left(t,z;x\right) e^{i\frac{\rho_n^2}{\alpha}\left(\alpha^2-\delta^2\right)x} e^{-\alpha\rho_n z} e^{\delta\rho_n t'} dt'. \qquad (3.3\text{c})
\end{aligned}
$$

The final solution is work on $z = t$. Hence, by substituting eq.$(3.3a)$ to eq.$(3.3b)$ and eq.$(3.3c)$ we find the solution (for $n = 1, 2, ..., N$):

$$
q_1 = \frac{-A_{n_0}\, e^{\rho_n(\delta-\alpha)t} e^{-i\frac{\rho_n^2}{\alpha}\left(\delta^2-\alpha^2\right)x}}{1 \,+\, \left(\frac{\mu\left(\left|A_{n_0}\right|^2+\left|B_{n_0}\right|^2\right)}{\left(\left(\frac{\alpha^2-\delta^2}{\alpha}\right)^{1/2}\left(\rho_n-\sigma_n\right)\right)^2}\right) e^{\rho_n(\delta-\alpha)t-i\frac{\rho_n^2}{\alpha}\left(\delta^2-\alpha^2\right)x} e^{-\sigma_n(\delta-\alpha)t+i\frac{\sigma_n^2}{\alpha}\left(\delta^2-\alpha^2\right)x}}\,, \qquad (3.4\text{a})
$$

and

$$
q_2 = \frac{-B_{n_0} e^{\rho_n(\delta-\alpha)t} e^{-i\frac{\rho_n^2}{\alpha}\left(\delta^2-\alpha^2\right)x}}{1 + \left(\frac{\mu\left(\left|A_{n_0}\right|^2+\left|B_{n_0}\right|^2\right)}{\left(\left(\frac{\alpha^2-\delta^2}{\alpha}\right)^{1/2}\left(\rho_n-\sigma_n\right)\right)^2}\right) e^{\rho_n(\delta-\alpha)t-i\frac{\rho_n^2}{\alpha}\left(\delta^2-\alpha^2\right)x} e^{-\sigma_n(\delta-\alpha)t+i\frac{\sigma_n^2}{\alpha}\left(\delta^2-\alpha^2\right)x}}. \qquad (3.4\text{b})
$$

We define

$$\eta_n = k_n \left(t + ik_n\xi x\right), \tag{3.5a}$$

and

$$\eta_n^* = k_n^* \left(t - ik_n^*\xi x\right) \quad , \tag{3.5b}$$

where

$$k_n = \left(\delta - \alpha\right)\rho_n \ , \tag{3.5c}$$

$$k_n^* = -\left(\delta - \alpha\right)\sigma_n \ , \tag{3.5d}$$

$$\tau = \left(\frac{\alpha^2 - \delta^2}{\alpha}\right)^{1/2}\left(\frac{1}{\delta - \alpha}\right), \tag{3.5e}$$

and

$$\xi = \frac{\left(\alpha^2 - \delta^2\right)}{\alpha\left(\delta - \alpha\right)^2}. \tag{3.5f}$$

Here ρ_n and σ_n are arbitrary complex parameters and $\rho_n^* = -\sigma_n$.

Now q_1 and q_2 can be rewritten as

$$q_1 = \sum_{n=1}^{N} \frac{-A_{n_0}\, e^{\eta_n}}{1 + e^{R_n + \eta_n + \eta_n^*}}, \tag{3.6a}$$

and

$$q_2 = \sum_{n=1}^{N} \frac{-B_{n_0}\, e^{\eta_n}}{1 + e^{R_n + \eta_n + \eta_n^*}}, \tag{3.6b}$$

where

$$e^{R_n} = \frac{\mu\left(\left|A_{n_0}\right|^2 + \left|B_{n_0}\right|^2\right)}{\left(\tau k_n + \tau k_n^*\right)^2}. \tag{3.7}$$

Based on our solution in eq.$(3.6a)$ and $(3.6b)$, we can see that our results are the bright and dark multi-soliton solution since $\mu = \pm\frac{\alpha^2-\delta^2}{\alpha^2\delta}$.The results of $q_1$and q_2 show that there are an collision of the bright and dark multi-soliton. If we choose parameter $\left(\left(\frac{(\alpha+\delta)}{\alpha(\alpha-\delta)}\right)\mu\right) > 0$ then we get the bright N-soliton solution. On the other hand, the dark N-soliton solution is found when $\left(\left(\frac{(\alpha+\delta)}{\alpha(\alpha-\delta)}\right)\mu\right) < 0$. In the equations, there are only four arbitrary complex parameters A_{n_0}, B_{n_0}, ρ_n and σ_n which can directly influence the phase of the solitons interaction. The results of q_1 and q_2 can also be rewritten in the more conventional form by introducing $\rho_n = l_n + i\lambda_n$ (where l_n and λ_n are arbitrary real parameters),

$$q_1 = \sum_{n=1}^{N} \frac{\left(\frac{-\left(\frac{\alpha^2-\delta^2}{\alpha}\right)^{1/2} l_n A_{n_0}}{\sqrt{\mu\left(\left|A_{n_0}\right|^2+\left|B_{n_0}\right|^2\right)}} \right) \exp\left(i\left(\delta-\alpha\right)\left[\lambda_n t + \xi\left(\delta-\alpha\right)\left(l_n^2-\lambda_n^2\right)x\right]\right)}{\cosh\left[\left(\delta-\alpha\right) l_n \left(t-2\xi\left(\delta-\alpha\right)\lambda_n x\right)+\varphi_n\right]}, \tag{3.8}$$

and

$$q_2 = \sum_{n=1}^{N} \frac{\left(\frac{-\left(\frac{\alpha^2-\delta^2}{\alpha}\right)^{1/2} l_n B_{n_0}}{\sqrt{\mu\left(\left|A_{n_0}\right|^2+\left|B_{n_0}\right|^2\right)}} \right) \exp\left(i\left(\delta-\alpha\right)\left[\lambda_n t + \xi\left(\delta-\alpha\right)\left(l_n^2-\lambda_n^2\right)x\right]\right)}{\cosh\left[\left(\delta-\alpha\right) l_n \left(t-2\xi\left(\delta-\alpha\right)\lambda_n x\right)+\varphi_n\right]}, \tag{3.9}$$

where φ_n is a real multi-soliton phase,

$$\varphi_n = \frac{1}{2} R_{n,} \tag{3.10}$$

and the amplitudes of the multi-soliton are

$$Amp^1 = -\frac{\left(\frac{\alpha^2-\delta^2}{\alpha}\right)^{1/2} l_n A_{n_0}}{\sqrt{\mu\left(\left|A_{n_0}\right|^2+\left|B_{n_0}\right|^2\right)}}, \tag{3.11a}$$

and

$$Amp^2 = -\frac{\left(\frac{\alpha^2-\delta^2}{\alpha}\right)^{1/2} l_n B_{n_0}}{\sqrt{\mu\left(\left|A_{n_0}\right|^2+\left|B_{n_0}\right|^2\right)}}. \tag{3.11b}$$

Parameter λ_n contributes to the velocities of the multi-soliton.[15]

The results derived above have contributed on the obtaining a general form of an exact real parameter μ and have also shown that there is a general multi-soliton solution of the integrable coupled **NLS** eq. of Manakov type (eq.(1.1)).

Based on our results in eq.(3.6a) and (3.6b), the bright N-solitons solution (for $\left(\left(\frac{(\alpha+\delta)}{\alpha(\alpha-\delta)}\right)\mu\right) > 0$) can be reduced to the bright one and two soliton solutions related to that in the works that have been done before by Radhakrishnan, et. al. using Hirota method in Ref.10. According to the comparison of both methods, their results of the bright one soliton solution is equal to our results when $\alpha\, e^{\eta_1^{(0)}} = -A_{1_0}$, $\beta e^{\eta_1^{(0)}} = -B_{1_0}$, $\frac{(\alpha+\delta)}{\alpha(\alpha-\delta)} = 1$, $\xi = 1$, $\tau = 1$, $k_1 = (\delta - \alpha)\,(l_1 + i\lambda_1), \mu = +\frac{\alpha^2-\delta^2}{\alpha^2\delta}$ and $(\mid \alpha \mid^2 + \mid \beta \mid^2) = \left(|A_{1_0}|^2 + |B_{1_0}|^2\right)$. On the other hand, their results of the inelastic collision of the bright two soliton can be reduced to our elastic collision of the solution if we put $\alpha_1 : \alpha_2 = \beta_1 : \beta_2$ in their result.

We can also compare our results with the results in Ref.8. We find that our result and the Shchesnovich result are the same when $\chi = \frac{1}{2}$, $2\mu = 1$, $x = z$, $t = \tau$, $n = 1$ and $|\theta_1|^2 + |\theta_2|^2 = |A_{1_0}|^2 + |B_{1_0}|^2 = 1$. So, we can generalize that our result is the solution of the bright and dark multi-soliton collisions.

IV. Discussions and Conclusions

We have presented a general form of the bright and dark multi-soliton solution of the integrable coupled NLS equation of Manakov type using the inverse scattering Zakharov-Shabat scheme.We can conclude that the solution of the Manakov type can be solved by using an expanded inverse scattering Zakharov-Shabat scheme in which we have chosen a certain operator in eq.(2.1), and a certain k_+ (eq.(2.7a)) and F (eq.(2.7b)) in our solution.

Finally, we find the novel result provided in Ref.9 and the results of the use of Hirota method in Ref.10 and Rev.11 that the solution in eq.(3.6a) and (3.6b) corresponds to an elastic collision of the bright and dark multi-soliton, as long as $A_{1_0} : A_{2_0} : ... : A_{N_0}$ is equal to $B_{1_0} : B_{2_0} : ... : B_{N_0}$. If in our results, $A_{1_0} : A_{2_0} : ... : A_{N_0} \neq B_{1_0} : B_{2_0} : ... : B_{N_0}$, then we get an inelastic collision of the bright and dark multi-solitons. From the eq.(3.8) and eq.(3.9), we can also conclude that although the collision between the bright and dark multi-solitons their velocities and amplitudes or intensities do not change, their phases do change.

According to the comparison between our bright one and two soliton solution of the integrable coupled **NLS** equation of Manakov type and the solution provided by Radhakrishnan, R., et.al., we get that their inelastic collision of the bright two-soliton solution can be our elastic collision solution, if some certain parameters in the results are omitted by putting $\alpha_1 : \alpha_2 = \beta_1 : \beta_2$. In our results, the parameter $\exp\left(\eta_j^{(0)}\right)$ appeared in Ref.10

has been absorbed into A_{n_0} and B_{n_0}. Based on our results, we propose that there must be an integrable m-tupled (vector) multi-solitons solution of the following equation:[4,16,17]

$$\left(i\frac{\partial}{\partial x}+\chi\frac{\partial^2}{\partial t^2}+2\mu\sum_{b=1}^{m}|q_b|^2\right)q_c \;=\; 0, \;\; c=1,2,...,m,$$

where χ and μ are arbitrary real parameters. We also propose that there must be an elastic and inelastic collision of the bright and dark multi-soliton in this system.[17]

ACKNOWLEDGEMENTS

Both authors would like to thank H.J.Wospakrik for useful discussions. We also thank Yu. S. Kivshar and N.N. Akhmediev (Optical Sciences Centre, The Institute for Advanced Studies, The Australian National University) for their information and suggestions. We also would like to thank P. Silaban for his encouragements. The work of F.Z. is supported by the Hibah Bersaing VII/1, 1998-1999, DIKTI, Republic of Indonesia.The work of H.E. is partially supported by CIDA - EIUDP, Republic of Indonesia.

References

[1] M. Remoissenet, *Waves Called Solitons; Concepts and Experiments*, Springer-Verlag, Berlin, and references therein (1994).

[2] R. Radhakrishnan and M. Lakshmanan, J. Phys. A **28**, 2683 (1995).

[3] R. Radhakrishnan, M. Lakshmanan, and J. Hietarinta, Phys. Rev. E. **56**, 2213 (1997).

[4] N.N. Akhmediev, W. Królikowski, and A.W. Snyder, Phys. Rev. Lett. **81**, 4632-4633 (1998).

[5] A.P. Seppard and Yu. S. Kivshar, Phys. Rev. E. **55**, 4773-4782 (1997).

[6] F.P. Zen and H.I. Elim, *Soliton Solution of the Integrable Coupled Nonlinear Schrödinger Equation of Manakov Type*, hep-th/9812215.

[7] V.S. Shchesnovich, and E.V. Doktorov, Phys. Rev. E. **55**,7626 (1997).

[8] V. Shchesnovich, *Polarization Scattering by Soliton-Soliton Collisions*, solv-int/ 9712020.

[9] H.J.S. Dorren, *On the Integrability of Nonlinear Partial Differential Equations*, solv-int/9807007.

[10] R. Radhakrishnan, M. Lakshmanan, and J. Hietarinta, *Inelastic Collision and Switching of Coupled Bright Solitons in Optical Fibers*, solv-int/9703008.

[11] R. Radhakrishnan and M. Lakshmanan, J. Phys. A ***28,*** 2683 (1995).

[12] M.J. Ablowitz and H. Segur, *Solitons and the inverse scattering transform*, SIAM studies in applied mathematics, Philadelpia, p.p.228-230 (1981).

[13] P.G. Drazin and R.S. Johnson, *Solitons: an Introduction*, Cambridge University Press, Cambridge, p.p.144-160 (1992).

[14] R.K. Bullough and P.J. Caudrey, *Topics in Current Physics*, Springer-Verlag, Berlin, p.p.253-268.

[15] N.N. Akhmediev and A. Ankiewicz, 1997, *Solitons; Nonlinear Pulses and Beams*, Chapman & Hall, London, p.p.66-70 (1980).

[16] L.D. Faddeev and L.A. Takhtajan, (1987), *Hamiltonian Methods in the Theory of Solitons*, Springer-Verlag, London, p.288.

[17] F.P. Zen and H.I. Elim, in preparation.